


Collisions at the Large Hadron Collider



Ritchie Patterson

CIPT
July 12, 2010



Dark Matter

Dark matter seems to cause the lensing arcs in this Hubble image




Massey et al (Caltech) used arcs like these to create a map of the dark matter in a section of the sky. (2007)

Dark Matter and the Universe

What do we know about it?

Not much.

- It exerts gravitational force – so it has mass
- It is widely distributed, but not uniformly (this rules out very light particles such as neutrinos).
- It rarely interacts with particles we know -- quarks, electrons, neutrinos, etc. (that's why we can't see it.)




It's not any of the particles we know.

- 13.7 billion years ago there was a giant explosion
- 10^{-20} seconds later, the universe was a hot soup of particles and radiation
- No planets, no stars, not even atoms
- Instead, particles and radiation in *thermal equilibrium* ($E=mc^2$)

- Then the universe cooled
- Most heavy particles decayed away, and radiation had too little energy to produce more.
- Atomic nuclei formed
- Then stars and galaxies
- And planets
- And you and me

- Some heavy particles may have remained as dark matter
- These may be recreated only when enough energy is available
- In nature
- Or at the Large Hadron Collider

Where are the gaps?






Some big gaps in understanding

- What is the dark matter?
- Where did the antimatter go?
- Why do particles have mass?
 - What is the Higgs particle and why is it so light?
- What is dark energy?
- What was the nature of gravity in the very early universe?

The LHC will push back to the energies of the early universe to address some of these questions


How do colliders work?


- They use $E = mc^2$
- Accelerate particles to near the speed of light and smash them together. Their kinetic energy and mass energy combine to create heavy particles.
- At the LHC, protons are accelerated and the quarks inside them collide. The total energy will be $E = 14 \text{ TeV}$.




What do you need?

- Lots of protons to smash together
 - Proton source (hydrogen tank)
- Mechanism for accelerating them
 - Protons at the LHC travel at 99.99% of the speed of light
- Racetrack (ring) for storing and colliding beams of protons
 - Only a couple of protons interact each time the beams collide, so we collide them over and over again.
- Detector to observe, and disentangle the collision products







Overall view of the LHC



The ring

Electric and magnetic fields
accelerate and steer the beams



- Beams are in 2808 bunches, each with 1.15×10^{11} protons
- Total energy in each beam is 360 MJ (equivalent to **60 kg TNT**)


LHC Steering Magnet

The last one being
lowered into place



The magnets

- Steer the protons around the 17 mile circular track
- Are superconducting with a field of 8.3 Tesla
- Use 96 tons of superfluid helium





Proton-proton collision




Particle detectors

- Heavy particles produced in the collisions typically decay within 10^{-24} seconds into particles we know and love: e, mu, pi, K
- The detector records traces of these well-known and well-understood particles.



CMS

ATLAS




What do we want from our detector?


Imagine that a bomb explodes mid-air, and you want to study the fragments to find out everything you can about the bomb.

What properties of the fragments would you want to measure?

- Direction of motion of each fragment just after explosion
- Speed (or momentum) of each fragment
- Mass of each fragment



Each particle species leaves a distinct trail




CMS Detector at the LHC


- CMS stands for “Compact Muon Solenoid”
- Weighs 12,500 tons
- Six stories tall

A detector cross-section, showing particle paths

- Beam Pipe (center)
- Tracking Chamber
- Magnet Coil
- E-M Calorimeter
- Hadron Calorimeter
- Magnetized Iron
- Muon Chambers



CMS
A Compact Solenoidal Detector for LHC



Total weight : 12,500t
Overall diameter : 13.00 m
Overall length : 21.60 m
Magnetic field : 4 Tesla



CMS-PARA-001-11/07/97

JLB,PP

CMS is “compact” because its diameter is 60% that of ATLAS, the other big LHC detector. But CMS weighs 1.7 times more.


CMS Tracker

- Solid state detector
- Charged particles produce a signal (or “hit”) on each layer
- Resolution is 10-60 microns per hit




Immersed in 4 Tesla
solenoidal magnetic field ...
Why?

Particles in the tracker



$B = 4$
T



Which particle(s)
have the greatest
momentum?



The smallest?

Which have positive
electric charge?

Electromagnetic Calorimeter

Absorbs electrons and photons to measure their energies

Simulated “shower”
in crystal




Lead-tungstate
crystals



Shower particles produce
scintillation light, detected at
the back of the crystal


Electromagnetic calorimeter



76,000
Lead-tungstate
crystals




Slice of CMS



Computing

- 15 Petabytes of data per year
- Tiered computing structure
- Achieved data transfer rates of 110 Gbps
- the world record




CMS Cavern

92 meters underground




Installing CMS




- Components were built on the surface and then lowered via crane.
- This is the largest one, weighing more than 2000 tons.
- After lowering, they move on air pads.



LHC timeline






CMS Experiment at LHC, CERN
Run 135149, Event 125426133
Lumi section: 1345
Sun May 09 2010, 05:24:09 CEST

$Z \rightarrow \mu\mu$

Muon $p_T = 67.3, 50.6 \text{ GeV}/c$
Inv. mass = $93.2 \text{ GeV}/c^2$








CMS Experiment at LHC, CERN
Run 133877, Event 28405693
Lumi section: 387
Sat Apr 24 2010, 14:00:54 CEST

Electrons $p_T = 34.0, 31.9 \text{ GeV}/c$
Inv. mass $= 91.2 \text{ GeV}/c^2$

$Z \rightarrow ee$








CMS Experiment at LHC, CERN
Run 133875, Event 1228182
Lumi section: 16
Sat Apr 24 2010, 09:08:46 CEST

Muon $p_T = 38.7 \text{ GeV}/c$
 $ME_T = 37.9 \text{ GeV}$
 $M_T = 75.3 \text{ GeV}/c^2$

$W \rightarrow \mu \nu$






CMS Experiment at LHC, CERN
Run 133874, Event 21466935
Lumi section: 301
Sat Apr 24 2010, 05:19:21 CEST

Electron $p_T = 35.6 \text{ GeV}/c$
 $ME_T = 36.9 \text{ GeV}$
 $M_T = 71.1 \text{ GeV}/c^2$

$W \rightarrow e \nu$




Results so far

- Momentum and angular distributions of particles produced in proton collisions
- Identification of familiar particles
- Results will probe deeper in the future:
 - With more collisions, you can see rarer, heavier particles. (Important for detecting higgs)
 - Better calibration of the detector allows more nuanced analysis. (Important for seeing dark matter).

Finding Dark Matter


- Leaves no trace in detector
- Find via momentum imbalance of detected particles: $\Sigma \mathbf{p} \neq 0$



This is a simulation of dark matter in ATLAS

This event is modeled based on a candidate theory called supersymmetry.

Other theories give similar momentum imbalance.



Conclusions

The LHC is now operating and CMS and ATLAS are working well.

First results have been published (more soon). More collisions and results will come over the next few years.

They will explore energies not seen since moments after the big bang.